For Research Use Only

CEP152 Polyclonal antibody

Catalog Number: 21815-1-AP

Featured Product

3 Publications



Purification Method:

WB 1:200-1:1000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

21815-1-AP NM_014985

Size: Genel D (NCBI): 150ul, Concentration: 350 µg/ml by 22995

Nanodrop and 287 µg/ml by Bradford Full Name:

method using BSA as the standard;

ethod using BSA as the standard; centrosomal protein 152kDa

Source: Calculated MW:
Rabbit 189 kDa
Isotype: Observed MW:
IgG 66 kDa

Positive Controls:

WB: mouse brain tissue,

IHC: human brain tissue,

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

IF. WB

Species Specificity:

human, mouse

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

CEP152, also named as KIAA0912, is a regulator of genomic integrity and cellular response to DNA damage acting through ATR-mediated checkpoint signaling. It is necessary for centrosome duplication. CEP152 functions as a molecular scaffold facilitating the interaction of PLK4 and CENPJ, two molecules involved in centriole formation. CEP152, ATR and CENPJ control cellular responses to DNA damage. CEP152 has some isoforms with MW 189 kDa, 152 kDa, 110 kDa and 66 kDa.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|------------|-------------|
| Zhenguang Ying | 36280838 | BMC Biol | WB,IF |
| Tianyu Wu | 36395215 | Science | IF |
| I-Ling Lu | 29950674 | Nat Commun | WB,IF |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 21815-1-AP (CEP152 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 21815-1-AP (CEP152 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



IP result of anti-CEP152(IP:21815-1-AP, 4ug; Detection:21815-1-AP 1:2000) with mouse brain tissue lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 21815-1-AP (CEP152 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).