For Research Use Only

FOXA2 Polyclonal antibody Catalog Number:22474-1-AP Featured Product

Featured Product 23 Publications




Basic Information	Catalog Number: 22474-1-AP	GenBank Accession Num BC011780	ber: Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 550 ug/ml by	3170	WB 1:500-1:2000	
	Nanodrop;	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source:	Q9Y261	protein lysate	
	Rabbit	Full Name:	IHC 1:2000-1:8000 IF-P 1:50-1:500	
	Isotype:	forkhead box A2		
	IgG	Calculated MW:		
	Immunogen Catalog Number: AG18221	457 aa, 48 kDa		
		Observed MW: 55-65 kDa		
Applications	Tested Applications: Positive Controls:			
	WB, IHC, IF-P, IP, ELISA	w	WB : HT-29 cells, HepG2 cells, A549 cells, MDA-MB-231	
	Cited Applications:	CE	ells	
	WB, IHC, IF, CoIP, chIP	IP	: HepG2 cells,	
	Species Specificity: human	IF	IC : human colon cancer tissue,	
	Cited Species:	IF	-P : rat stomach tissue,	
	human, mouse			
	Note-IHC: suggested antigen (TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
	bujjer pri 0.0			
Background Information	FOXA2 also named as HNF 3 beta or domains. FOXA2 can Shuttle betwee FOXA2 Thr-156 abolishes binding to FOXA2 as a transcription factor invol	n the nucleus and cytoplas target promoters and subso ves in embryonic develop pression in differentiated	m in a CRM1-dependent manner. Phosphorylation of	
	FOXA2 also named as HNF 3 beta or domains. FOXA2 can Shuttle betwee FOXA2 Thr-156 abolishes binding to FOXA2 as a transcription factor invol expression and regulation of gene ex but the phosphorylated FOXA2 may b	n the nucleus and cytoplas target promoters and subso ves in embryonic develop pression in differentiated		
	FOXA2 also named as HNF 3 beta or domains. FOXA2 can Shuttle betwee FOXA2 Thr-156 abolishes binding to FOXA2 as a transcription factor invol expression and regulation of gene ex but the phosphorylated FOXA2 may be Author Pub	n the nucleus and cytoplas target promoters and subse ves in embryonic develops pression in differentiated pe a 55-65kDa protein.	m in a CRM1-dependent manner. Phosphorylation of equent transcription activation upon INS stimulation ment, establishment of tissue-specific gene tissues. The molecular weight of FOXA2 is 48kDa, Application	
Background Information Notable Publications	FOXA2 also named as HNF 3 beta or domains. FOXA2 can Shuttle betwee FOXA2 Thr-156 abolishes binding to FOXA2 as a transcription factor invol expression and regulation of gene ex but the phosphorylated FOXA2 may be Author Pub Lin Yang 317	n the nucleus and cytoplas target promoters and subse- ves in embryonic develop pression in differentiated be a 55-65kDa protein. med ID Journal	m in a CRM1-dependent manner. Phosphorylation or equent transcription activation upon INS stimulation nent, establishment of tissue-specific gene tissues. The molecular weight of FOXA2 is 48kDa, Application Sci WB	
	FOXA2 also named as HNF 3 beta or domains. FOXA2 can Shuttle betwee FOXA2 Thr-156 abolishes binding to FOXA2 as a transcription factor invol expression and regulation of gene exputs but the phosphorylated FOXA2 may beAuthorPubLin Yang317 Yan HuangYan Huang363	n the nucleus and cytoplas target promoters and subse ves in embryonic develops pression in differentiated be a 55-65kDa protein. med ID Journal 54329 Int J Biol S	m in a CRM1-dependent manner. Phosphorylation of equent transcription activation upon INS stimulation nent, establishment of tissue-specific gene tissues. The molecular weight of FOXA2 is 48kDa, Application Sci WB als WB,IF	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 22474-1-AP (FOXA2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 22474-1-AP (FOXA2 antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 22474-1-AP (FOXA2 antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-FOXA2 (IP:22474-1-AP, 4ug; Detection:22474-1-AP 1:500) with HepG2 cells lysate 1320 ug.



Immunofluorescent analysis of (4% PFA) fixed rat stomach tissue using FOXA2 antibody (22474-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).