## For Research Use Only

## HOXA7 Polyclonal antibody

Catalog Number:22931-1-AP



**Basic Information** 

Catalog Number:

22931-1-AP

Nanodrop:

GenBank Accession Number:

BC148692

GeneID (NCBI):

150ul, Concentration: 350 ug/ml by 3204

**UNIPROT ID:** 

P31268

Rabbit Full Name:

Isotype: homeobox A7

IgG Calculated MW: Immunogen Catalog Number: 230 aa, 25 kDa

Observed MW:

22 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

AG18927

Species Specificity:

human

Positive Controls:

WB: HEK-293T cells, COLO 205 cells, THP-1 cells

**Purification Method:** 

WB 1:500-1:2000

Antigen Affinity purified

Recommended Dilutions:

## **Background Information**

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 22931-1-AP (HOXA7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.