

For Research Use Only

FIZ1 Polyclonal antibody

Catalog Number: 23257-1-AP



Basic Information

Catalog Number: 23257-1-AP	GenBank Accession Number: BC136512	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 587 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 84922	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	UNIPROT ID: Q96SL8	
Isotype: IgG	Full Name: FLT3-interacting zinc finger 1	
Immunogen Catalog Number: AG19760	Calculated MW: 496 aa, 52 kDa	
	Observed MW: 52 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Jurkat cells, NIH/3T3 cells, HEK-293 cells
Species Specificity: human, mouse	

Background Information

FLT3 interacting zinc finger 1 (FIZ1) is a ubiquitous expressed protein that composes of 11 C2H2 zinc finger domains spread over the entire protein in clusters of two, three, or four fingers. The interaction of FIZ1 was mapped to the catalytic domain of FLT3, which is a receptor tyrosine kinase involved in proliferation and survival of hematopoietic progenitor cells as well as differentiation of early B lymphoid progenitors (10409713). In the mammalian retina, FIZ1 interacts with NRL (Neural-Retina Leucine-zipper), an essential transcriptional activator of rod photoreceptor-specific genes, and the concentration of FIZ1 in the retina will increase during photoreceptor terminal maturation (18854042).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 23257-1-AP (FIZ1 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.