For Research Use Only

IL-2 Polyclonal antibody Catalog Number: 26156-1-AP 15 Publications




Basic Information	Catalog Number: 26156-1-AP	GenBank Accession Numb NM-008366	er: Purification Method: Antigen affinity purification
	Size:	GeneID (NCBI):	Recommended Dilutions:
	150ul , Concentration: 1000 ug/ml by Nanodrop and 333 ug/ml by Bradford		WB 1:500-1:2000 IHC 1:50-1:500
	method using BSA as the standard;	Full Name: interleukin 2	1.50-1.500
	Source: Rabbit	Calculated MW: 19 kDa	
	lsotype: IgG		
	Immunogen Catalog Number: AG24079		
Applications	Tested Applications:	Positive Controls:	
	Cited Applications: cells,		: PMA, Ionomycin and Brefeldin A treated EL-4
			HC : human liver cancer tissue, human tonsillitis
	Species Specificity: human, mouse	tissue	
	Cited Species: human, mouse, rat		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w	vely, antigen	
	buffer pH 6.0		
Background Information	Interleukin-2 (IL2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tra tumors. Nevertheless, it has been sho	ces the proliferation of thes eat immune-deficiencies lik own that Il2 also acts as the tions in the target cell throu	te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (Il2R). Il2R is a
	Interleukin-2 (IL2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tra tumors. Nevertheless, it has been sho cells. Interleukin 2 mediates its funct multimeric functional protein consist	ces the proliferation of thes eat immune-deficiencies lik own that Il2 also acts as the tions in the target cell throu	e and others cells like CD8+ T cells, B and NK cell te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (Il2R). Il2R is a
	Interleukin-2 (II2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tra tumors. Nevertheless, it has been sho cells. Interleukin 2 mediates its funct multimeric functional protein consist Author Pu	ces the proliferation of thes eat immune-deficiencies lik wm that IL2 also acts as the tions in the target cell through ting of three different chains	e and others cells like CD8+ T cells, B and NK cell te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (Il2R). Il2R is a t: Il2Rα, Il2Rβ and γ c.
	Interleukin-2 (IL2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tre tumors. Nevertheless, it has been sho cells. Interleukin 2 mediates its funct multimeric functional protein consist Author Pu Abdolrazagh Hashemi Shahraki 35	ces the proliferation of thes eat immune-deficiencies lik won that IL2 also acts as the tions in the target cell throu- ting of three different chains ubmed ID Journal	e and others cells like CD8+ T cells, B and NK cell te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (II2R). II2R is a the interleu
	Interleukin-2 (II2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tra tumors. Nevertheless, it has been sho cells. Interleukin 2 mediates its funct multimeric functional protein consist Author Pu Abdolrazagh Hashemi Shahraki 35 Dongxu Lin 35	ces the proliferation of thes eat immune-deficiencies lik own that IL2 also acts as the tions in the target cell through ting of three different chains ibmed ID Journal 5717488 Lung	e and others cells like CD8+ T cells, B and NK cell te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (II2R). II2R is a the interleukin 2 receptor (II2R). II2R is a till2R0, II2Rβ and γ c. Application WB ol Res WB
Background Information Notable Publications	Interleukin-2 (II2) is a protein initiall CD4+CD25- (helper) T cells, and indu For this reason, it has been used to tra tumors. Nevertheless, it has been sho cells. Interleukin 2 mediates its funct multimeric functional protein consist Author Pu Abdolrazagh Hashemi Shahraki 35 Dongxu Lin 35	ces the proliferation of thes eat immune-deficiencies lik with that IL2 also acts as the tions in the target cell through ing of three different chains ibmed ID Journal i717488 Lung i755168 J Immund i2426338 Lipids He er shipment.	e and others cells like CD8+ T cells, B and NK cell te HIV, and induce immune response against main growth factor of CD4+CD25+ Regulatory T gh the interleukin 2 receptor (II2R). II2R is a the interleukin 2 receptor (II2R). II2R is a till2R0, II2Rβ and γ c. Application WB ol Res WB

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





PMA, A23187 and Brefeldin A treated EL-4 cells were subjected to SDS PAGE followed by western blot with 26156-1-AP (IL-2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 26156-1-AP (II2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 26156-1-AP (Il2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).