

For Research Use Only

BCMA/TNFRSF17 Polyclonal antibody, PBS Only

Catalog Number: 27724-1-PBS



Basic Information

Catalog Number:

27724-1-PBS

Size:

100ug, Concentration: 1 mg/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG27049

GenBank Accession Number:

BC058291

GeneID (NCBI):

608

UNIPROT ID:

Q02223

Full Name:

tumor necrosis factor receptor
superfamily, member 17

Calculated MW:

184 aa, 20 kDa

Observed MW:

20 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human

Background Information

BCMA (B cell maturation antigen), also known as TNFRSF17, is 20.2-kDa type III transmembrane glycoprotein and is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes and plasma cells, which may be important for B cell development and autoimmune response (PMID: 32943087; 9846698). BCMA has two agonist ligands: a proliferation-inducing ligand (APRIL) and BAFF. When BCMA binds to APRIL, it transmits signals of cell survival and proliferation (PMID: 27127303); when BCMA binds to BAFF, it mediates the activation of NF- κ B and MAPK8/JNK (PMID: 36140254). It has been found that the overexpression and activation of BCMA are associated with multiple myeloma (MM) in preclinical models and humans, supporting its potential utility as a therapeutic target for MM (PMID: 32055000; 32943087).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 27724-1-AP (TNFRSF17 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 27724-1-PBS in a different storage buffer formulation.



Daudi and Ramos cells were subjected to SDS PAGE followed by western blot with 27724-1-AP (TNFRSF17 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 27724-1-PBS in a different storage buffer formulation.