### For Research Use Only

# ICAM-1/CD54 Polyclonal antibody

30806-1-AP

Nanodrop:

Size:

IgG

Catalog Number: 30806-1-AP



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC015969

GeneID (NCBI):

150ul, Concentration: 500 ug/ml by 3383

Genein (NCBI):

ENSEMBL Gene ID:

Source: ENSG0000090339
Rabbit UNIPROT ID:
Isotype: P05362

Full Name:

Calculated MW: 90 kDa Observed MW:

85-95 kDa

intercellular adhesion molecule 1

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Ramos cells, LO2 cells, Raji cells

IHC: human ovary cancer tissue, human intrahepatic

**Purification Method:** 

WB 1:1000-1:8000 IHC 1:200-1:800

IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

cholangiocarcinoma tissue

IF/ICC: Raji cells,

## **Background Information**

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF-a, INF-y, IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).

### Storage

**Applications** 

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 30806-1-AP (ICAM-1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 30806-1-AP (ICAM-1/CD54 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 30806-1-AP (ICAM-1/CD54 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Raji cells using ICAM-1 antibody (30806-1-AP) at dilution of 1:400 and CoraLite\$488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (5A00013-2).