For Research Use Only

## EIF3F Polyclonal antibody Catalog Number: 31630-1-AP




Catalog Number: 31630-1-AP	GenBank Accession Number: BC000490	Purification Method: Antigen affinity Purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG34849	GenelD (NCBI): 8665	Recommended Dilutions: WB 1:5000-1:50000
	UNIPROT ID: O 00303	
	Full Name: eukaryotic translation initiation factor 3, subunit F Calculated MW: 38 kDa	
	Tested Applications:	Positive Co
WB: HeLa C		ells, Jurkat cells, mouse liver tissue, rat e
cells and is an integral part of the tra	nslation initiation factor 3(Eif3) con	plex. EIF3 is a complex of at least 10
Storage Buffer:		
	31630-1-AP Size: 150ul, Concentration: 500 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG34849 Tested Applications: WB, ELISA Species Specificity: human, mouse, rat EIF3F (eukaryotic translation initiatic cells and is an integral part of the tra- distinct subunits and is the largest tra- Storage: Storage Buffer: PBS with 0.02% sodium azide and 50	31630-1-AP BC000490   Size: GeneID (NCBI):   150ul, Concentration: 500 ug/ml by 8665   Nanodrop; UNIPROT ID:   Source: O00303   Rabbit Full Name:   Isotype: eukaryotic translation initiation   IgG factor 3, subunit F   Immunogen Catalog Number: Calculated MW:   AG34849 38 kDa   Observed MW: 40-45 kDa   Tested Applications: Positive Co   WB, ELISA WB : HeLa or   Species Specificity: brain tissue   human, mouse, rat EIF3F (eukaryotic translation initiation factor 3 subunit F) is a protein that cells and is an integral part of the translation initiation factor in eukaryot   Storage: Storage:   Storage: Storage is translation on year after shipment.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31630-1-AP (EIF3F antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.