For Research Use Only

Ki-67 Polyclonal antibody

Catalog Number: 32270-1-AP



Basic Information

Catalog Number:

32270-1-AP

Nanodrop:

GenBank Accession Number:

NM 002417

GeneID (NCBI): Size: 150ul, Concentration: 450 ug/ml by 4288

UNIPROT ID: Source: P46013 Rabbit Full Name:

Isotype: antigen identified by monoclonal

antibody Ki-67 IgG

> Calculated MW: 359 kDa

Applications

Tested Applications:

IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue, human colon cancer

Purification Method:

IHC 1:500-1:2000 IF/ICC 1:200-1:800

Antigen affinity Purification

Recommended Dilutions:

tissue, human ovary cancer tissue

IF/ICC: HeLa cells,

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 32270-1-AP (Ki-67 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 32270-1-AP (Ki-67 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Ki-67 antibody (32270-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4), CL488-phalloidin (green).