For Research Use Only

p150 glued Polyclonal antibody


Catalog Number:55182-1-AP 2 Publications



Basic Information	Catalog Number: 55182-1-AP	GenBank Accession Number: NM_004082	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 500 µg/ml by	1639	WB 1:500-1:2000	
	Nanodrop and 167 µg/ml by Bradford method using BSA as the standard; Source:	Full Name: dynactin 1 (p150, glued homolo Drosophila)	IHC 1:20-1:200 pg, IF 1:20-1:200	
	Rabbit	Calculated MW:		
	Isotype:	142 kDa		
	IgG	Observed MW: 135 kDa, 150 kDa		
Applications	Tested Applications:	Positive Controls:		
	IF, IHC, WB,ELISA Cited Applications:	WB : human brain tissue, Jurkat cells		
	IF, WB		human brain tissue,	
	Species Specificity: IF : MDCK cells, human, Canine			
	Cited Species: human, mussel			
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen		
	DCTN1, also named as DAP-150, p150-glued and p135, belongs to the dynactin 150 kDa subunit family. It is required for the cytoplasmic dynein-driven retrograde movement of vesicles and organelles along microtubules. Dynein- dynactin interaction is a key component of the mechanism of axonal transport of vesicles and organelles. Defects ir DCTN1 are the cause of distal hereditary motor neuronopathy type 7B (HMN7B). This antibody is specific to DCTN1.			
Background Information	dynactin interaction is a key component	ent of the mechanism of axonal		
	dynactin interaction is a key compon DCTN1 are the cause of distal heredit	ent of the mechanism of axonal		
	dynactin interaction is a key component of the cause of distal heredit Author Public	ent of the mechanism of axonal ary motor neuronopathy type 7E	3 (HMN7B). This antibody is specific to DCTN1	
	dynactin interaction is a key component DCTN1 are the cause of distal heredit Author Pub Tianyu Wu 363	ent of the mechanism of axonal any motor neuronopathy type 7E med ID Journal	3 (HMN7B). This antibody is specific to DCTN1 Application IF	
Background Information Notable Publications Storage	dynactin interaction is a key compone DCTN1 are the cause of distal heredit Author Pub Tianyu Wu 363 Jian-Wei Zheng 300 Storage: Stora at -20°C. Storage Buffer: PBS with 0.02% sodium azide and 50°	ent of the mechanism of axonal ary motor neuronopathy type 7E med ID Journal 95215 Science 07186 Ecotoxicol Envi % glycerol pH 7.3.	3 (HMN7B). This antibody is specific to DCTN1 Application IF	
Notable Publications	dynactin interaction is a key compone DCTN1 are the cause of distal heredit Author Pub Tianyu Wu 363 Jian-Wei Zheng 300 Storage: Stora at -20°C. Storage Buffer:	ent of the mechanism of axonal ary motor neuronopathy type 7E med ID Journal 95215 Science 07186 Ecotoxicol Envi % glycerol pH 7.3.	3 (HMN7B). This antibody is specific to DCTN1 Application IF	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human brain tissue were subjected to SDS PAGE followed by western blot with 55182-1-AP (DCTN1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain using 55182-1-AP (DCTN1 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain using 55182-1-AP (DCTN1 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed MDCK cells using 55182-1-AP (p150 glued antibody) at dilution of 1:50 and Alexa Fluor 488conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).