For Research Use Only

6*His, His-tag Monoclonal antibody

Catalog Number:60001-1-lg 5 Publications



Basic Information

Catalog Number:

60001-1-lg

150ul , Concentration: 200 $\mu g/ml$ by Bradford method using BSA as the standard;

Source:

Mouse Isotype:

IgG2b

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Calculated MW:

Purification Method:

Recommended Dilutions:

WB 1:200-1:1000

Applications

Tested Applications:

WB. FIISA

Cited Applications:

Species Specificity: recombinant protein

Positive Controls:

WB: Recombinant Protein,

Background Information

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His tags are most often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized $metal\ ions, including\ nickel, cobalt\ and\ copper, under\ specific\ buffer\ conditions.\ As\ is\ the\ case\ with\ other\ protein\ tag$ systems, this polyhistidine tag can often be cleaved at sites recognized by proteases such as thrombin and enterokinases to isolate the protein of interest. The His tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Wei-Dong Zhao	29895952	Nat Commun	
Ying Ji	29901114	Oncol Rep	WB
Shu-Hong Zhang	30792309	J Biol Chem	

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data