

For Research Use Only

GAPDH Monoclonal antibody

Catalog Number: 60004-1-Ig **6000 Publications**



Basic Information

Catalog Number: 60004-1-Ig	GenBank Accession Number: BC004109	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 2597	CloneNo.: 1E6D9
Source: Mouse	Full Name: glyceraldehyde-3-phosphate dehydrogenase	Recommended Dilutions: WB 1:50000-1:500000 IP 0.5-4.0 ug for IP and 1:2000-1:12000 for WB
Isotype: IgG2b	Calculated MW: 36 kDa	IF 1:200-1:2000
Immunogen Catalog Number: AG0766	Observed MW: 36 kDa	

Applications

Tested Applications: FC, IF, IP, WB, ELISA	Positive Controls:
Cited Applications: ChIP, ColP, FC, IF, IHC, IP, WB	WB : HeLa cells, HepG2 cells, ROS1728 cells, pig brain tissue, zebrafish tissue, whole yeast, whole Nematode tissue, soybean whole plant tissue, arabidopsis whole plant tissue, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, C6 cells, PC-12 cells, C2C12 cells, SP2/0 cells, rat brain tissue, mouse brain tissue
Species Specificity: human, mouse, rat, yeast, plant, zebrafish	IP : HeLa Cells,
Cited Species: Arabidopsis, Bombyx mori, bovine, Branchiostoma belcheri, brovine, Caenorhabditis elegans, canine, chicken, CHO cell line, Cynomorium songaricum	IF : Ethacrynic acid treated HeLa cells,

Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types. For murine tissue samples, conjugated mouse antibody HRP-60004 and rabbit antibody 10494-1-AP are preferable.

Notable Publications

Author	Pubmed ID	Journal	Application
Shenli Zhang	34600025	Food Chem Toxicol	WB
Yanliang Wu	34601083	J Ethnopharmacol	WB
Jiangang Zheng	34587973	BMC Vet Res	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:200000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.




Western blot analysis of GAPDH in various tissues and cell lines using Proteintech antibody 60004-1-Ig at a dilution of 1:10000.


IP & WB of 60004-1-Ig with HeLa Cell




IP result of anti-GAPDH (60004-1-Ig for IP and Detection) with HeLa cell lysate.



Arabidopsis, rice, wheat, corn whole plant tissue were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH Antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of EA treated HeLa cells using 60004-1-Ig (GAPDH antibody) at dilution of 1:50 and Rhodamine-labeled goat anti-mouse IgG (red).



1X10⁶ HeLa cells were stained with 0.2ug GAPDH antibody (60004-1-Ig, red) and control antibody (60004-1-Ig, blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). FITC-Goat anti-Mouse IgG with dilution 1:100.