

For Research Use Only

GAPDH Monoclonal antibody, PBS Only



Catalog Number: 60004-1-PBS

Basic Information

Catalog Number: 60004-1-PBS	GenBank Accession Number: BC004109	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 2597	CloneNo.: 1E6D9
Source: Mouse	UNIPROT ID: P04406	
Isotype: IgG2b	Full Name: glyceraldehyde-3-phosphate dehydrogenase	
Immunogen Catalog Number: AG0766	Calculated MW: 36 kDa	
	Observed MW: 36 kDa	

Applications

Tested Applications:
WB, IP, IF, FC, ELISA

Species Specificity:
human, mouse, rat, yeast, plant, zebrafish

Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types. For murine tissue samples, conjugated mouse antibody HRP-60004 and rabbit antibody 10494-1-AP are preferable.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.


Storage Buffer:
PBS only

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:200000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.



Western blot analysis of GAPDH in various tissues and cell lines using Proteintech antibody 60004-1-Ig at a dilution of 1:10000. This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.





Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.


IP & WB of 60004-1-Ig with HeLa Cell



Arabidopsis, rice, wheat, corn whole plant tissue were subjected to SDS PAGE followed by western blot with 60004-1-Ig (GAPDH Antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of EA treated HeLa cells using 60004-1-Ig(GAPDH antibody) at dilution of 1:50 and Rhodamine-labeled goat anti-mouse IgG (red). This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer formulation.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human GAPDH (60004-1-Ig, Clone:1E6D9) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 60004-1-PBS in a different storage buffer