## For Research Use Only

## IL-18 Monoclonal antibody, PBS Only

Catalog Number:60070-1-PBS Featured Product



**Basic Information** 

Catalog Number:

GenBank Accession Number:

BC007461

**Purification Method:** Protein A purification

60070-1-PBS

GeneID (NCBI):

CloneNo.:

100ug, Concentration: 1mg/ml by

5C6F8

Nanodrop;

**UNIPROT ID:** Q14116 Full Name:

Mouse Isotype:

IL-18 Calculated MW:

Immunogen Catalog Number:

193 aa, 22 kDa

AG1063

lgG2b

Observed MW:

24 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, FC (Intra), ELISA

Species Specificity:

human

**Background Information** 


 $IL18, is a proinflam matory\ cytokine\ involved\ in\ the\ development\ of\ Th1\ cells\ and\ in\ immune\ response.\ It\ can$ stimulate the NK cells and certain T cells to release IFN gamma which plays an important role in activating the macrophages or other cells. IL18 has been demonstrated to have the potential to enhance Fas ligand-mediated cytotoxicity, which is increased in PE and regulates placental apoptosis. IL18 is synthesized as a 24 kDa precursor and then cleaved into a biologically activate 18 kDa form.

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60070-1-lg (IL18 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.




WB result of Pre-IL18 antibody (60070-1-1g, 1:4000) with si-Control and si-Pre-IL18 transfected HeLa cells.. This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60070-1-lg (IL18 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60070-1-lg (IL18 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60070-1-1g (IL-18 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-18 antibody (60070-1-lg, Clone: 5C6F8) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using IL-18 antibody (60070-1-Ig, Clone: 5C6F8) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (5A00013-1). This data was developed using the same antibody clone with 60070-1-PBS in a different storage buffer formulation.



1x10^6 HeLa cells were intracellularly stained with 0.4 ug IL-18 Monoclonal antibody (60070-1-Ig. Clone:5C6F8) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.4 ug Mouse IgG2b isotype control Mouse McAb (66360-3-Ig, Clone: 11B8C4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 60070-1-PBS in a

