For Research Use Only

PAX8 Monoclonal antibody, PBS Only

Catalog Number:60145-4-PBS




Basic Information	Catalog Number: 60145-4-PBS	GenBank Accession Number: BC001060	Purification Method: Protein A purification
	Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG2b Immunogen Catalog Number: AG0306	GeneID (NCBI): 7849 UNIPROT ID: Q06710 Full Name: paired box 8 Calculated MW: 48 kDa Observed MW: 48 kDa	CloneNo.: 4H7B3
Applications	Tested Applications: WB, IHC, IF/ICC, IF-P, Indirect ELISA Species Specificity: human, mouse, rat		
Background Information	PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX8 has 5 isoforms with calculated MW 31kd, 34kd, 41kd, 43kd and 48kd. And there is a 60kd band corresponding to the electrophoretic mobility of PAX8 (PMID:15650356).		
Storage	Storage: Store at -80°C. Storage Buffer: PBS Only		


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60145-4-Ig (PAX8 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60145-4-Ig (PAX8 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation.



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 60145-4-1g (PAX8 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation.



Confocal immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using 60145-4-Ig (PAX8 antibody) at dilution of 1:100 and Coralite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Phalloidin stains F-actin (Red). This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation. Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 60145-4-1g (PAX8 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse kidney tissue using PAX8 antibody (60145-4-1g, Clone: 4H7B3) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Mouse $\lg(H+L)$ (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60145-4-PBS in a different storage buffer formulation.