

For Research Use Only

NCAM1/CD56 Monoclonal antibody, PBS Only

Catalog Number:60238-1-PBS



Basic Information

Catalog Number: 60238-1-PBS	GenBank Accession Number: BC047244	Purification Method: Protein A purification
Size: 100ug , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 4684	CloneNo.: 1E8C9
Source: Mouse	ENSEMBL Gene ID: ENSG00000149294	
Isotype: IgG1	UNIPROT ID: P13591	
Immunogen Catalog Number: AG5732	Full Name: neural cell adhesion molecule 1	
	Calculated MW: 95 kDa	
	Observed MW: 140 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, IF-P, ELISA

Species Specificity:
human, rat, pig

Background Information

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronectin-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isoforms have also been described (PMID: 1856291).


Storage

Storage:
Store at -80°C.
Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




fetal human brain tissue were subjected to SDS PAGE followed by western blot with 60238-1-Ig (NCAM1/CD56 antibody at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human colon tissue using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60238-1-PBS in a different storage buffer formulation.