

For Research Use Only

IL-19 Monoclonal antibody, PBS Only



Catalog Number: 60278-1-PBS

Basic Information

Catalog Number: 60278-1-PBS	GenBank Accession Number: BC153127	Purification Method: Protein G purification
Size: 100ug, Concentration: 1mg/ml by Nanodrop;	GeneID (NCBI): 29949	CloneNo.: 8H10H8
Source: Mouse	UNIPROT ID: Q9UHD0	
Isotype: IgG1	Full Name: interleukin 19	
Immunogen Catalog Number: AG18887	Calculated MW: 177 aa, 20 kDa	

Applications

Tested Applications:
IHC, IF/ICC, IF-P, Indirect ELISA

Species Specificity:
human

Background Information

IL-19, a member of the IL-10 cytokine family, is expressed in epithelial cells, endothelial cells, and macrophages. It can bind the interleukin-20 receptor complex IL-20R1/IL-20R2 and lead to the activation of the signal transducer and activator of transcription 3 (STAT3). IL-19 induces IL-6 and TNF-alpha production in monocytes. It also induces cell apoptosis and reactive oxygen species production in monocytes, which suggests a role of this cytokine in inflammatory responses. It is up-regulated in monocytes following stimulation with lipopolysaccharide (LPS), and granulocyte-macrophage colony-stimulating factor (GM-CSF). IL-19 alters the balance of Th1 and Th2 cells in favor of Th2 cells. IL-19 contributes to a range of diseases and disorders, such as breast cancer, squamous cell carcinoma of the skin, asthma, endotoxic shock, uremia, psoriasis, rheumatoid arthritis, and periodontal and vascular disease.

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60278-1-Ig (IL-19 antibody at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60278-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60278-1-Ig (IL-19 antibody at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60278-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-19 antibody (60278-1-Ig, Clone: 8H10H8) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60278-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-19 antibody (60278-1-Ig, Clone: 8H10H8) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60278-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed U-937 cells using IL-19 antibody (60278-1-Ig, Clone: 8H10H8) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60278-1-PBS in a different storage buffer formulation.