For Research Use Only

## U2AF35 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

2B7F11

Catalog Number: 60289-1-PBS

**Basic Information** 

Catalog Number: GenBank Accession Number:

60289-1-PBS BC001923

Size: GeneID (NCBI): 100ug , Concentration: 1mg/ml by 7307

Nanodrop; UNIPROT ID:
Source: Q01081
Mouse Full Name:

Isotype: U2 small nuclear RNA auxiliary factor

IgG2c

Immunogen Catalog Number: Calculated MW: AG0399 28 kDa

Observed MW: 35-40 kDa

**Applications** 

**Tested Applications:** 

Indirect ELISA, IP, IF/ICC, IHC, WB

Species Specificity: rat, mouse, human


## **Background Information**

U2 auxiliary factor (U2AF), comprising a large and a small subunit, is a non-snRNP protein required for the binding of U2 snRNP to the pre-mRNA branch site. U2 (RNU2) small nuclear RNA auxiliary factor 1 (U2AF1,synonyms: RN, FP793, U2AF35, U2AFBP, RNU2AF1) is the small subunit which plays a critical role in both constitutive and enhancer-dependent RNA splicing by directly mediating interactions between the large subunit and proteins bound to the enhancers. The U2AF1 gene is localized to chromosome 21q22.3, which is the critical region for three diseases, progressive myoclonus epilepsy, autoimmune polyglandular disease type 1, and one form of bipolar affective disorder. The calculated molecular weight of U2AF35 is 28 kDa, but the modified U2AF35 protein is about 35-40 kDa.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## Selected Validation Data




WB result of U2AF35 antibody (60289-1-lg; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-U2AF35 transfected HeLa cells. This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60289-1-1g (U2AF35 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60289-1-lg (U2AF35 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.




IP result of anti-U2AF35 (IP:60289-1-Ig, 3ug; Detection:60289-1-Ig 1:500) with HeLa cells lysate 3000ug. This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using U2AF35 antibody (60289-1-Ig, Clone: 2B7F11) at dilution of 1:2000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using U2AF35 antibody (60289-1-lg, Clone: 2B7F11) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+I). This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.



1X10^6 Ramos cells were intracellularly stained with 0.4 ug Anti-Human U2AF35 (60289-1-lg, Clone:2B7F11) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 60289-1-PBS in a different storage buffer formulation.