For Research Use Only

Perilipin-2 Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

1G5D4

Catalog Number: 60340-1-PBS

Basic Information

Catalog Number:

60340-1-PBS

Nanodrop:

Mouse

GenBank Accession Number:

BC005127

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

UNIPROT ID: Q99541 Full Name:

Isotype: adipose differentiation-related

lgG1 protein

Immunogen Catalog Number: Calculated MW:

48 kDa AG7643

> Observed MW: 47-52 kDa

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT $family\ which\ is\ responsible\ for\ the\ transportation\ of\ lipids\ and\ the\ formation\ of\ lipid\ droplets.\ ADRP\ is\ localized\ on$ the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage

Storage: Store at -80°C. Storage Buffer:

PBS Only


Selected Validation Data



human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-lg (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-lg (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat brown adipose slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-lg (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.