For Research Use Only

Anti-Human CD3 (Hit3a)

Catalog Number:65112-1-Ig



Purification Method:

Affinity purification

CloneNo.:

Hit3a

Basic Information

Catalog Number:

65112-1-lg

Size:

500ug , 0.5 mg/ml

Source: Mouse Isotype: IgG2a, kappa GenBank Accession Number:

BC049847 GeneID (NCBI):

916

ENSEMBL Gene ID: ENSG00000198851

UNIPROT ID:

P07766 Full Name:

CD3e molecule, epsilon (CD3-TCR

complex)
Calculated MW:
207 aa, 23 kDa

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65112-1-1g, Clone:Hit3a) or Mouse IgG2a Isotype Control and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500, then CL750 Anti-Human CD45. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65112-1-1g, Clone:Hit3a) or Mouse IgGZa Isotype Control and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500. Cells were not fixed. Lymphocytes were gated.