

Anti-Human CD154 (24-31)

Catalog Number: 65156-1-Ig

Basic Information

Catalog Number: 65156-1-Ig	GenBank Accession Number: BC071754	Purification Method: Purified by protein-A affinity chromatography
Size: 100ug, 1 mg/ml	GeneID (NCBI): 959	CloneNo.: 24-31
Source: Mouse	Full Name: CD40 ligand	
Isotype: IgG1, kappa	Calculated MW: 261 aa, 29 kDa	

Applications

Tested Applications:
FC

Species Specificity:
Human, Non-human primates

Background Information

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4⁺ T cells and on a small proportion of CD8⁺ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.

Storage


Storage:
Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.1% sodium azide.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ PMA (50 ng/ml) and ionomycin (1 ug/ml)-stimulated human PBMCs (5 hours) were surface stained with 0.2 ug Anti-Human CD154 (65156-1-Ig, Clone: 24-31) and FITC anti-mouse IgG1 Antibody at dilution 1:200 (green), or stained with 0.2 ug isotype control and FITC anti-mouse IgG1 Antibody at dilution 1:200 (black). Cells were not fixed.