

For Research Use Only

Anti-Human M-CSF (3G12E6)

Catalog Number: 65283-1-Ig



Basic Information

Catalog Number: 65283-1-Ig	GenBank Accession Number: BC021117	Purification Method: Affinity purification
Size: 100ug, 1000 µg/ml	GeneID (NCBI): 1435	CloneNo.: 3G12E6
Source: Mouse	Full Name: colony stimulating factor 1 (macrophage)	
Isotype: IgG1	Calculated MW: 60 kDa	
Immunogen Catalog Number: HZ-1192		

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human

Background Information

M-CSF plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes. M-CSF promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. On the cellular level, M-CSF promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration and plays a role in lipoprotein clearance (PMID: 29734839; 29679908; 28202039)

Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ LPS and TNF- α then Brefeldin A treated human PBMCs were surface stained with CoraLite[®] Plus 405 Anti-Human CD11b then intracellularly stained with 0.2 ug CoraLite[®] Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated. This data was developed using the CoraLite[®] Plus 488 conjugated antibody



1X10⁶ LPS and TNF- α then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite[®] Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite[®] Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit