For Research Use Only

Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody

Catalog Number: 65559-1-MR



Basic Information

Catalog Number:

65559-1-MR

Size:

100ug , 1 mg/ml

Source: Mouse Isotype: IgG2a

GenBank Accession Number: BC025715

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000153563

UNIPROT ID: P01732

Full Name: CD8a molecule Calculated MW: 235 aa, 26 kDa

Purification Method: Protein A purfication

CloneNo.: UCHT4

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody (65559-1-MR, Clone: UCHT4) or Mouse IgG2a Isotype Control, and PEconjugated Goat Anti-House IgG. Cells were costained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (APC-65569, Clone: OKT3). Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody (65559-1-MR, Clone: UCHT4) (red) or Mouse IgG2a Isotype Control, and PE-conjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated.