For Research Use Only

Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

10.1

Catalog Number: 65572-1-PBS

Basic Information

Catalog Number: 65572-1-PBS

1mg, 2 mg/ml

Mouse Isotype:

Size:

Source:

lgG2a

ENSEMBL Gene ID: ENSG00000150337

GeneID (NCBI):

BC032634

2209

Full Name:

Fc fragment of IgG, high affinity Ia,

GenBank Accession Number:

receptor (CD64) Calculated MW: 374 aa, 43 kDa

Applications

Tested Applications:

Species Specificity:

Human

Background Information

Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\).$ CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.


Storage

Store at -80°C. Stable for one year after shipment.

Storage Buffer: PBS Only

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.5 ug Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody, PBS Only (65572-1-PBS, Clone:10.1) or 0.5 ug Mouse IgG2a Isotype Control (C1.18.4), (65208-1-Ig, Clone: C1.18.4), CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), and 0.5 ug APC Anti-Human CD11b (ICRF44) (APC-65116, Clone: ICRF44). Cells were not fixed. Monocytes were gated.