For Research Use Only

Anti-Human CD86 (BU63) Mouse IgG2a Recombinant Antibody

Catalog Number: 65592-1-MR



Purification Method:

CloneNo.:

BU63

Protein A purification

Basic Information

Catalog Number:

65592-1-MR

Size:

100ug , 1 mg/ml

Source: Mouse Isotype: lgG2a

ENSEMBL Gene ID: ENSG00000114013 Full Name:

GeneID (NCBI):

BC040261

GenBank Accession Number:

CD86 molecule Calculated MW: 329 aa, 38 kDa

Applications

Tested Applications:

Species Specificity:

Human

Background Information

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding $of CD86\ with\ CD28\ antigen\ is\ a\ costimulatory\ signal\ for\ activation\ of\ the\ T-cell.\ Binding\ of\ CD86\ with\ cytotoxic\ T-cell.$ $lymphocyte-associated\ protein\ 4\ negatively\ regulates\ T-cell\ activation\ and\ diminishes\ the\ immune\ response.$


Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD86 Mouse IgG2a Recombinant Antibody (65592-1-MR, Clone: BU63) or Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) and Coralite® 488-Conjugated Donkey Anti-Mouse IgG(H+L). Cells were then stained with APC Anti-Human CD14. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD86 Mouse IgG2a Recombinant Antibody (65592-1-MR, Clone:BU63) (red) or Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue) and Coralite® 488-Conjugated Donkey Anti-Mouse IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.