For Research Use Only

Anti-Human CD27 (O323) Mouse IgG2a Recombinant Antibody

Catalog Number: 65627-1-MR



Purification Method:

CloneNo.:

0323

Protein A purification

Basic Information

Catalog Number:

65627-1-MR

Size:

100ug , 1 mg/ml

Source: Mouse

Isotype: IgG2a GenBank Accession Number:

BC012160 GeneID (NCBI):

939

ENSEMBL Gene ID:

ENSG00000139193 Full Name:

CD27 molecule
Calculated MW:

29 kDa

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD27 (0323) Mouse IgG2a Recombinant Antibody (65627-1-MR, Clone:0323) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4), and PE-conjugated Goat Anti-Mouse IgG. Cells were then stained with APC Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD27 (0323) Mouse IgG2a Recombinant Antibody (65627-1-MR, Clone: 0323) (red) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue), and PEconjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated.