For Research Use Only

Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody

Catalog Number: 65630-1-MR



Purification Method:

Protein A purification

CloneNo.:

SK1

Basic Information

Catalog Number:

65630-1-MR

Size:

Source:

100ug , 1 mg/ml

Mouse Isotype: lgG2a

GenBank Accession Number:

BC025715

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000153563

CD8a molecule Calculated MW:

235 aa, 26 kDa

Full Name:

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody (65630-1-MR, Clone: SK1) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4), and PE-conjugated Goat Anti-Mouse IgG. Cells were then stained with APC Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody (65630-1-MR, Clone: SK1) (red) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue), and PEconjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated.