

For Research Use Only

RBAP48 Monoclonal antibody, PBS Only



Catalog Number: 66060-1-PBS

Featured Product

Basic Information

Catalog Number:

66060-1-PBS

Size:

100ug, Concentration: 1mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG6196

GenBank Accession Number:

BC053904

GeneID (NCBI):

5928

UNIPROT ID:

Q09028

Full Name:

retinoblastoma binding protein 4

Calculated MW:

48 kDa

Observed MW:

53 kDa

Purification Method:

Protein A purification

CloneNo.:

5C4D6

Applications

Tested Applications:

Indirect ELISA, IF/ICC, IHC, WB

Species Specificity:

rat, mouse, human

Background Information

Histone-binding protein RBBP4 (also known as RbAp48, or NURF55) is a protein that in humans is encoded by the RBBP4 gene. This gene encodes a ubiquitously expressed nuclear protein that belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2/NuRD complex complex that has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of corepressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. A decrease of RbAp48 in the dentate gyrus (DG) of the hippocampus in the brain is suspected to be a main cause of memory loss in normal aging (PMID: 23986399).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66060-1-Ig (RBAP48 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 66060-1-Ig (RBAP48 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human cervical cancer using 66060-1-Ig (RBAP48 antibody) at dilution of 1:200 (under 10x lens). This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 66060-1-Ig (RBAP48 antibody) at dilution of 1:200 (under 40x lens). This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer formulation.



1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug Anti-Human RBAP48 (66060-1-Ig, Clone:5C4D6) and CoraLite[®]488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer



WB result of RBAP48 antibody (66060-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RBAP48 transfected HeLa cells. This data was developed using the same antibody clone with 66060-1-PBS in a different storage buffer formulation.