For Research Use Only

## S100A6 Monoclonal antibody, PBS Only

Catalog Number:66098-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

66098-1-PBS

GeneID (NCBI):

BC001431

Protein A purification

100ug, Concentration: 1mg/ml by

CloneNo.: 3D11C8

Nanodrop:

**UNIPROT ID:** P06703

Mouse

Full Name:

Isotype: lgG1

S100 calcium binding protein A6 Calculated MW:

Immunogen Catalog Number:

10 kDa

AG17606

Observed MW:

10 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human

**Background Information** 


S100A6 is also named as calcyclin, prolactin receptor-associated protein (PRA), growth factor-inducible protein 2A9 ir MLN4. It belongs to \$100 family of low molecular weight, acidic, calcium-binding proteins which contain two EFhand calcium binding sites. S100A6 may function as a calcium sensor to activate several processes in the calcium signal transduction pathway of cell growth, proliferation, secretion and exocytosis. Although S100A6 is a cytoplasmic protein, it can be located in the nuclear and cytoplasmic membranes in the presence of Ca2+ (PMID: 37670392).

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only


## Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 66098-1-1g (\$100A6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66098-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66098-1-lg (S100A6 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66098-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66098-1-lg (S100A6 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66098-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 66098-1-lg (\$100A6 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66098-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using \$100A6 antibody (66098-1-Ig, Clone: 3D11C8) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66098-1-PBS in a different storage buffer formulation.