

For Research Use Only

Prealbumin/transthyretin Monoclonal antibody, PBS Only



Catalog Number: 66108-1-PBS

Basic Information

Catalog Number: 66108-1-PBS	GenBank Accession Number: BC020791	Purification Method: Protein A purification
Size: 100ug , Concentration: 1mg/ml by Nanodrop;	GeneID (NCBI): 7276	CloneNo.: 1H8C4
Source: Mouse	UNIPROT ID: P02766	
Isotype: IgG2a	Full Name: transthyretin	
Immunogen Catalog Number: AG19232	Calculated MW: 147 aa, 16 kDa	
	Observed MW: 14-16 kDa, 28-33 kDa	

Applications

Tested Applications:
Indirect ELISA, IHC, WB

Species Specificity:
mouse, human

Background Information

Transthyretin (TTR) is a plasma transport protein for thyroxine and retinol, through the association with retinol-binding protein. It is a homotetrameric protein synthesized mainly in liver, choroid plexus, retinal pigment epithelium, and pancreas. Within the CNS, TTR is the only known protein synthesized solely by the choroid plexus. Mutant and wildtype TTR give rise to various forms of amyloid deposition (amyloidosis). Defects in TTR are the cause of amyloidosis transthyretin-related (AMYL-TTR), hyperthyroxinemia dystransthyretinemic euthyroidal (HTDE) and carpal tunnel syndrome type 1 (CTS1). In addition, positive immunostaining for TTR has been reported as a sensitive diagnostic marker of choroid plexus tumors. TTR can be expressed as monomer (14-16 kDa), dimer (28-33 kDa), or tetramer around 55-60 kDa. (PMID: 22103483)

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 66108-1-Ig (Prealbumin/transferrin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66108-1-PBS in a different storage buffer formulation.



1×10^6 PC-12 cells were intracellularly stained with 0.4 ug Anti-Human Prealbumin/transferrin (66108-1-Ig, Clone:1H8C4) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66108-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66108-1-Ig (Prealbumin/transferrin antibody at dilution of 1:50. This data was developed using the same antibody clone with 66108-1-PBS in a different storage buffer formulation.