

For Research Use Only

FABP5 Monoclonal antibody, PBS Only



Catalog Number: 66299-1-PBS

Featured Product

Basic Information

Catalog Number:

66299-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG3005

GenBank Accession Number:

BC019385

GeneID (NCBI):

2171

UNIPROT ID:

Q01469

Full Name:

fatty acid binding protein 5 (psoriasis-associated)

Calculated MW:

135 aa, 15 kDa

Observed MW:

15 kDa

Purification Method:

Protein A purification

CloneNo.:

1C6E12

Applications

Tested Applications:

WB, IF, IHC, ELISA

Species Specificity:

human, mouse, rat

Background Information

FABP5, also named as PA-FABP and E-FABP, belongs to the calycin superfamily and Fatty-acid binding protein (FABP) family. It is high specificity for fatty acids. FABP5 is highest affinity for C18 chain length. It may be involved in keratinocyte differentiation. FABP5 is a fatty acid-binding protein and is expressed in epidermis and endothelial cells of the microvasculature of different organs. FABP5 has also been identified as a tumor-associated antigen, which is highly expressed in various cancers. FABP5 was detected in the sera of HNSCC patients with early stage cancer. Antibodies specific for FABP5 were significantly increased in a substantial amount in patients, suggesting that FABP5 may be a potential diagnostic biomarker for HNSCC. FABP5 may serve as a biomarker for HNSCC. (PMID:19602232)

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.




WB result of FABP5 antibody (66299-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FABP5 transfected A375 cells. This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using FABP5 antibody (66299-1-Ig, Clone: 1C6E12) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.




A375 cells were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 Antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.





Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66299-1-PBS in a different storage buffer formulation.