## For Research Use Only

## MMP3 Monoclonal antibody

Catalog Number:66338-1-lg Featured Product

36 Publications




Basic Information	Catalog Number: 66338-1-lg	GenBank Accession Number: BC074869		Purification Metho Protein A purificati			
	Size:	GeneID (NCBI): 4314 UNIPROT ID: P08254 Full Name: matrix metallopeptidase 3 (stromelysin 1, progelatinase)		CloneNo.: 1F5A9			
	150ul, Concentration: 1000 µg/ml by						
	Bradford method using BSA as the standard;			Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000			
	Source:						
	Mouse						
	Isotype: IgG1 Immunogen Catalog Number: AG12359						
		Calculated MW:					
		477 aa, 54 kDa					
		Observed MW: 45-60 kDa					
Applications	Tested Applications: WB, IHC, ELISA		Positive Controls:				
				WB: PC-12 cells, human heart tissue, HeLa cells, feta			
	Cited Applications: WB, IF, IHC			human brain tissue, COLO 320 cells, Caco-2 cells, p heart tissue, pig brain tissue, rat brain tissue, mouse brain tissue			
	Species Specificity:		brain tissue				
	human, rat, mouse, pig		IHC : human breast cancer tissue, human colon ca				
	Cited Species: human, rat, mouse, pig				ontissue		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen					
			Matrix metalloproteinases (MMPs) play a critically important role in extracellular matrix remodeling and have bee implicated in a number of key normal and pathologic processes. These proteases have come to represent important therapeutic and diagnostic targets for the treatment and detection of human cancers. MMP-3 activate procollagenas via two pathways: slow direct activation and rapid activation in conjunction with tissue or plasma proteinases. The pro-MMP3 (60 kDa) and the active MMP3 (47 kDa) can be detected through western blot.				
Background Information	implicated in a number of key norma therapeutic and diagnostic targets for via two pathways: slow direct activat	l and pathologic proce r the treatment and de tion and rapid activation	sses.These prote tection of huma on in conjunctio	eases have come to n cancers. MMP-3 act n with tissue or plası	represent importai ivate procollagena		
	implicated in a number of key norma therapeutic and diagnostic targets for via two pathways: slow direct activat pro-MMP3 (60 kDa) and the active MM	l and pathologic proce r the treatment and de tion and rapid activation	sses.These prote tection of huma on in conjunction tected through v	eases have come to n cancers. MMP-3 act n with tissue or plası	represent importar ivate procollagena		
Background Information	implicated in a number of key norma therapeutic and diagnostic targets for via two pathways: slow direct activat pro-MMP3 (60 kDa) and the active MM Author Pub	l and pathologic proce r the treatment and de tion and rapid activati AP3 (47 kDa) can be de med ID Journ	sses.These prote tection of huma on in conjunction tected through v	eases have come to n cancers. MMP-3 act n with tissue or plası	represent importar ivate procollagena na proteinases. Th		
	implicated in a number of key norma therapeutic and diagnostic targets for via two pathways: slow direct activat pro-MMP3 (60 kDa) and the active MM Author Pub Wenbin Pei 346	l and pathologic proce r the treatment and de tion and rapid activation MP3 (47 kDa) can be de med ID Journ 550433 Front	sses.These prot tection of huma on in conjunctio tected through v	eases have come to n cancers. MMP-3 act n with tissue or plası	represent importan ivate procollagena na proteinases. Th Application		
	implicated in a number of key norma therapeutic and diagnostic targets for via two pathways: slow direct activat pro-MMP3 (60 kDa) and the active MM Author Pub Wenbin Pei 346 Weichao Sun 361	l and pathologic proce r the treatment and de tion and rapid activation (A7 kDa) can be de med ID Journ (50433 Front (02306 Int J	sses.These prot tection of human on in conjunction tected through v nal Pharmacol	eases have come to n cancers. MMP-3 act n with tissue or plası	represent importan ivate procollagena na proteinases. Th Application WB		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66338-1-1g (MMP3 antibody) at dilution of 1:4000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66338-1-1g (MMP3 antibody) at dilution of 1:4000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66338-1-lg (MMP3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of MMP3 antibody (66338-1-Ig; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-MMP3 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 66338-1-lg (MMP3 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).