

For Research Use Only

HVCN1 Monoclonal antibody, PBS Only



Catalog Number: 66449-1-PBS

Basic Information

| | | |
|---|---|---|
| Catalog Number: 66449-1-PBS | GenBank Accession Number: BC032672 | Purification Method: Protein G purification |
| Size: 100ug , Concentration: 1 mg/ml by Nanodrop; | GeneID (NCBI): 84329 | CloneNo.: 1E4C4 |
| Source: Mouse | UNIPROT ID: Q96D96 | |
| Isotype: IgG1 | Full Name: hydrogen voltage-gated channel 1 | |
| Immunogen Catalog Number: AG5350 | Calculated MW: 273 aa, 32 kDa | |
| | Observed MW: 28-35 kDa, 40 kDa, 60 kDa | |

Applications

Tested Applications:
WB, IF, IHC, Indirect ELISA

Species Specificity:
human

Background Information

HVCN1, also named as VSOP and HV1, Belongs to the hydrogen channel family. HVCN1 mediates the voltage-dependent proton permeability of excitable membranes. It forms a proton-selective channel through which protons may pass in accordance with their electrochemical gradient. Proton efflux, HVCN1 is accompanied by membrane depolarization, facilitates acute production of reactive oxygen species in phagocytosis. HVCN1, the voltage-sensitive proton channel, is present in human sperm and is an important regulator of the functional maturation of sperm. HVCN1 has four isoforms with MW 28-32 kDa or 40 kDa (modification). It has a dimer form with MW 60 kDa.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 66449-1-Ig (HVCN1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66449-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-Ig, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66449-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-Ig, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66449-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66449-1-Ig (HVCN1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66449-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66449-1-Ig (HVCN1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66449-1-PBS in a different storage buffer formulation.