For Research Use Only

CRABP2 Monoclonal antibody, PBS Only

Catalog Number:66468-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66468-1-PBS

BC001109 GeneID (NCBI): Protein A purification

Size:

by 1382

CloneNo.: 1A5F3

100ug, Concentration: 1mg/ml by Nanodrop:

UNIPROT ID: P29373

Source: Mouse

Full Name:

Isotype:

cellular retinoic acid binding protein

lgG1

2

Immunogen Catalog Number: AG0309

Calculated MW: 16 kDa

Observed MW:

14 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig


Background Information

Cellular retinoic acid binding protein 2 (CRABP2, synonyms: RBP6, CRABP-II). A number of specific carrier proteins for members of the vitamin A family have been discovered. Cellular retinoic acid binding proteins (CRABP) are low molecular weight proteins whose precise function remains unknown. CRABP2 is important in retinoic acid-mediated regulation of human skin growth and differentiation. It has been postulated that the CRABP2 gene is transcriptionally regulated by a newly synthesized regulatory protein.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




T-47D cells were subjected to SDS PAGE followed by western blot with 66468-1-Ig (CRABP2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




WB result of CRABP2 antibody (66468-1-Ig; 1:500; incubated at room temperature for 1.5 hours) with sh-Control and sh-CRABP2 transfected HEK-293 cells. This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human skin cancer tissue using CRABP2 antibody (66468-1-1g, Clone: 1A5F3) at dilution of 1:100 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+I). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66468-1-lg (CRABP2 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66468-1-lg (CRABP2 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CRABP2 antibody (66468-1-lg, Clone: 1A5F3) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(I+L) (SA00013-1). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human CRABP2 (66468-1-lg, Clone:1A5F3) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66468-1-