For Research Use Only

TGN46 Monoclonal antibody

Catalog Number:66477-1-lg 7 Publications



Basic Information

Catalog Number: GenBank Accession Number:

66477-1-lg BC028219 GeneID (NCBI): CloneNo.: Size: 150ul, Concentration: 1000 µg/ml by 10618 1F6D5

Source: trans-golgi network protein 2

Mouse Calculated MW: Isotype: 447 aa, 47 kDa lgG1 Observed MW: Immunogen Catalog Number: 90 kDa

AG4470

Applications

Tested Applications:

IHC, WB, ELISA **Species Specificity:** Human, Rat, Pig, Mouse

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein G purification

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:150-1:600

Positive Controls:

WB: A549 cells, HeLa cells, LNCaP cells, HepG2 cells, LO2 cells, SMMC-7721 cells, human colostrum tissue,

pig liver tissue

IHC: human breast cancer tissue, human liver cancer

tissue

Background Information

TGN46 (TGOLN2), the human homologue of rat Tgn38, is a transmembrane glycoprotein predominantly localized to the TGN (trans-Golgi network). TGN is a major secretory pathway sorting station for proteins and lipids. TGN46 may be involved in regulating membrane traffic to and from TGN. Alternatively spliced transcript variants encode different TGN46 isoforms. TGN46 has an apparent molecular mass of 100-150 kDa, suggesting extensive O- and Nglycosylations.

Notable Publications

Author **Pubmed ID** Application Haoyu Zhao 38230579 J Cell Physiol

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 66477-1-1g (TGN46 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66477-1-1g (TGN46 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66477-1-Ig (TGN46 antibody) at dilution of 1:300 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66477-1-lg (TGN46 antibody) at dilution of 1:300 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).