

For Research Use Only

HRPT2, CDC73 Monoclonal antibody, PBS Only



Catalog Number: 66490-1-PBS

Featured Product

Basic Information

Catalog Number:

66490-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG2959

GenBank Accession Number:

BC014351

GeneID (NCBI):

79577

UNIPROT ID:

Q6P1J9

Full Name:

cell division cycle 73, Paf1/RNA polymerase II complex component, homolog (S. cerevisiae)

Calculated MW:

531 aa, 61 kDa

Observed MW:

61 kDa

Purification Method:

Protein A purification

CloneNo.:

4B4H2

Applications

Tested Applications:

WB, IF, IHC, ELISA

Species Specificity:

Human, Pig, Mouse, Rat

Background Information

Parafibromin is a product of the hyperparathyroidism-jaw tumor syndrome gene HRPT2/CDC73, a putative tumor suppressor gene recently implicated in the autosomal dominant hyperparathyroidism-jaw tumor familial cancer syndrome, sporadic parathyroid cancer, and a minority of families with isolated hyperparathyroidism (PMID: 15580289). Defects in CDC73 are causes of hyperparathyroidism type 1 (HRPT1) and hyperparathyroidism type 2 (HRPT2) as well as parathyroid carcinoma (PRTC) (PMID: 12434154). Tumor suppressor parafibromin to the transcription elongation and RNA processing pathway as a PAF1 complex- and RNA polymerase II-bound protein (PMID: 15923622). Besides, parafibromin is also involved in post-transcriptional control pathways (PMID: 16116486). Recent report has revealed that pathogenic mutation, such as CDC73 gene mutation, is able to affect histone monoubiquitination (PMID: 22021426).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of HRPT2, CDC73 antibody (66490-1-Ig; 1:50000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HRPT2, CDC73 transfected HeLa cells. This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66490-1-Ig (HRPT2, CDC73 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66490-1-Ig (HRPT2, CDC73 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using HRPT2, CDC73 antibody (66490-1-Ig, Clone: 4B4H2) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66490-1-Ig (HRPT2, CDC73 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66490-1-Ig (HRPT2, CDC73 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66490-1-Ig (HRPT2, CDC73 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 66490-1-PBS in a different storage buffer formulation.