

For Research Use Only

# CCAR2 Monoclonal antibody, PBS Only



Catalog Number: 66497-1-PBS

Featured Product

## Basic Information

Catalog Number:

66497-1-PBS

Size:

100ug, Concentration: 1mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG18374

GenBank Accession Number:

BC018269

GeneID (NCBI):

57805

UNIPROT ID:

Q8N163

Full Name:

KIAA1967

Calculated MW:

923 aa, 103 kDa

Observed MW:

130 kDa

Purification Method:

Protein A purification

CloneNo.:

2D10F8

## Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, pig

## Background Information

Cell cycle and apoptosis regulator protein 2 (CCAR2), previously known as DBC1/KIAA1967, regulates diverse cellular functions including transcription, mRNA splicing, metabolism, and circadian cycle. CCAR2 is a critical regulator of cell death or survival following cellular stresses. CCAR2 is a widely expressed protein involved in the regulation of a variety of transcriptional complexes. High expression of CCAR2 correlates with poor outcomes in many human tumor types such as squamous cell carcinoma (SCC).

## Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66497-1-Ig (CCAR2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer formulation.




WB result of CCAR2 antibody (66497-1-Ig; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCAR2 transfected HEK-293 cells. This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66497-1-Ig (CCAR2 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66497-1-Ig (CCAR2 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 66497-1-Ig (CCAR2 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer formulation.



1X10<sup>6</sup> U2OS cells were intracellularly stained with 0.2 ug Anti-Human CCAR2 (66497-1-Ig, Clone:2D10F8) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66497-1-PBS in a different storage buffer