

For Research Use Only

TMEM173/STING Monoclonal antibody



Catalog Number: 66680-1-Ig

Featured Product

9 Publications

Basic Information

Catalog Number: 66680-1-Ig	GenBank Accession Number: BC047779	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 340061	CloneNo.: 1F1E1
Source: Mouse	Full Name: transmembrane protein 173	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:500-1:2000
Isotype: IgG2b	Calculated MW: 379 aa, 42 kDa	
Immunogen Catalog Number: AG20363	Observed MW: 37 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IF, IP, WB

Species Specificity:

Human, Rat

Cited Species:

human, mouse

Positive Controls:

WB : HSC-T6 cells, THP-1 cells, HEK-293 cells, HepG2 cells, Jurkat cells

IHC : human tonsillitis tissue, human spleen tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Stimulator of interferon genes (STING, also known as ERIS, MITA and MPYS, and encoded by TMEM173) is a transmembrane adaptor protein that facilitates innate immune signaling (PMID: 18724357). STING is widely expressed in various cell types such as endothelial cells, epithelial cells, T cells, macrophages, and dendritic cells (PMID: 26603901). It is predominantly located in the endoplasmic reticulum (ER). STING functions as a sensor of cytosolic DNA and promotes the production of type I interferons and pro-inflammatory cytokines. STING is a 379 amino acid protein with a calculated molecular weight of 42 kDa. It has been observed at 35-40 kDa (PMID: 27324217; 29632140; 30918080), and 70-80 kDa corresponding to the expected size of a STING dimer (PMID: 25790474; 29491158).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiaqi Shi	34182334	Cell Immunol	WB
Miao Zhao	33762743	Cell Death Differ	WB
Ziyang Feng	35248033	Cancer Cell Int	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HSC-T6 cells were subjected to SDS PAGE followed by western blot with 66680-1-Ig (TMEM173 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of TMEM173/STING antibody (66680-1-Ig; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-TMEM173/STING transfected THP-1 cells.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66680-1-Ig (TMEM173/STING antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66680-1-Ig (TMEM173/STING antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).