

For Research Use Only

# VWF Monoclonal antibody

Catalog Number: 66682-1-Ig **15 Publications**



## Basic Information

<b>Catalog Number:</b> 66682-1-Ig	<b>GenBank Accession Number:</b> GeneID (NCBI): 7450	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 150ul , Concentration: 1000 ug/ml by Nanodrop and 501 ug/ml by Bradford method using BSA as the standard;	<b>Full Name:</b> von Willebrand factor	<b>CloneNo.:</b> 3F9F3
<b>Source:</b> Mouse		<b>Recommended Dilutions:</b> IHC 1:250-1:1000 IF-P 1:200-1:800
<b>Isotype:</b> IgG1		
<b>Immunogen Catalog Number:</b> AG25578		

## Applications

<b>Tested Applications:</b> IHC, IF-P, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> IHC, IF	<b>IHC :</b> human breast cancer tissue, human liver cancer tissue, human prostate cancer tissue, human tonsillitis tissue
<b>Species Specificity:</b> human	<b>IF-P :</b> human breast cancer tissue,
<b>Cited Species:</b> human, mouse, rat	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

Von Willebrand factor (VWF) is a large multimeric glycoprotein found in blood plasma involved in hemostasis following vascular injury. Due to the multimeric nature of VWF, it can range in size from 500 to 20,000 kDa due to the differences in the number of subunits comprising the protein. Each subunit is approximately 250 kDa (PMID: 9759493). The biosynthesis of VWF in vivo is limited to endothelial cells (PMID: 4209883) and megakaryocytes (PMID: 2413071). VWF synthesized in endothelial cells is either released directly into the plasma via 27186a secretory pathway, or tubulized and stored in organelles unique to this cell type called Weibel-Palade bodies (PMID: 16459301). Whereas VWF synthesized in megakaryocytes is stored in the alpha granules of platelets (PMID: 2046403). The primary function of VWF is as an adhesive plasma glycoprotein, particularly factor VIII; an essential blood-clotting protein (PMID: 6982084). VWF is also important in platelet adhesion to wound sites by binding specifically to type I and type III collagen (PMID: 11098050), with larger VWF multimers being most effective (PMID: 24448155).

## Notable Publications

Author	Pubmed ID	Journal	Application
Lauren Mastrogiacomio	36499109	Int J Mol Sci	IF
Chaowei Hu	33215878	J Cell Mol Med	IHC
Weiqi Wu	35222704	Exp Ther Med	IF

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66682-1-Ig (vwf antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66682-1-Ig (vwf antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human breast cancer tissue using VWF antibody (66682-1-Ig, Clone: 3F9F3 ) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human breast cancer tissue using VWF antibody (66682-1-Ig, Clone: 3F9F3 ) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66682-1-Ig (VWF antibody) at dilution of 1:2000 (under 10x lens). Proteolytic pre-treatment mediated antigen retrieval with .