For Research Use Only

## ICOSLG/B7-H2/CD275 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

2G1F8

Catalog Number: 67003-1-PBS

**Basic Information** 

Catalog Number: GenBank Accession Number:

67003-1-PBS

100ug, Concentration: 1 mg/ml by

Nanodrop: **UNIPROT ID:** 075144 Mouse Full Name:

Isotype: inducible T-cell co-stimulator ligand IgG2a

BC064637

23308

GeneID (NCBI):

Calculated MW: Immunogen Catalog Number: 33 kDa, 34 kDa AG6799 Observed MW:

60-65 kDa, 28-35 kDa

**Applications** 

**Tested Applications:** WB, Indirect ELISA Species Specificity:

Human

**Background Information** 

ICOSLG, also known as B7H2 or GL50, is a type I transmembrane glycoprotein belonging to the B7 ligand family. ICOSLG is extensively expressed on professional antigen-presenting cells including B cells, macrophages, and dendritic cells, as well as non-lymphoid cells including mesenchymal stem cells, endothelial cells, fibroblasts, and tumor cells (PMID: 33756276; 35800767). It is a specific ligand for the T-cell-specific cell surface receptor ICOS and acts as a co-stimulatory molecule (PMID: 11023515; 11007762). The interaction of ICOSLG and ICOS plays important roles in the activation, proliferation, differentiation, and cytokine production of T cells as well as in the antibody secretion from B cells during secondary immune responses (PMID: 21851236).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67003-1-lg (ICOSLG antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67003-1-PBS in a different storage buffer formulation.