

For Research Use Only

# COMMD5 Monoclonal antibody, PBS Only



Catalog Number: 67043-1-PBS

## Basic Information

<b>Catalog Number:</b> 67043-1-PBS	<b>GenBank Accession Number:</b> BC003055	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug , Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 28991	<b>CloneNo.:</b> 1E10G6
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q9GZQ3	
<b>Isotype:</b> IgG2a	<b>Full Name:</b> COMM domain containing 5	
<b>Immunogen Catalog Number:</b> AG18464	<b>Calculated MW:</b> 25 kDa	
	<b>Observed MW:</b> 15 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
Human, Rat

## Background Information

COMM domain containing 5 (COMMD5, synonyms: HCARG, HT002) is a hypertension-related calcium-regulated gene. COMMD5 is negatively regulated by extracellular calcium concentration, and its basal mRNA levels were higher in hypertensive animals. Tissue distribution of COMMD5 shows a preponderance in the heart, stomach, jejunum, kidney (tubular fraction), liver, and adrenal gland (mainly in the medulla). COMMD5 mRNA is significantly more expressed in adult than in fetal organs, and its levels are decreased in tumors and cancerous cell lines. COMMD5 protein has no transmembrane domain, but contains 67% -helix content, a calcium-binding site, four putative "leucine zipper" motifs, and a nuclear receptor-binding domain. At the subcellular level, COMMD5 shows a nuclear localization.

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)      W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



human heart tissue were subjected to SDS PAGE followed by western blot with 67043-1-Ig (COMMD5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67043-1-PBS in a different storage buffer formulation.