## For Research Use Only

## ASC/TMS1 Monoclonal antibody

Catalog Number:67494-1-lg 30 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

67494-1-lg BC004470
Size: GeneID (NCBI):

150ul , Concentration: 1600 ug/ml by 29108
Nanodrop and 1000 ug/ml by Bradford\_UNIPROT ID:
method using BSA as the standard; Q9ULZ3

Source: Full Name:

Mouse PYD and CARD domain containing

Isotype:Calculated MW:IgG125 kDaImmunogen Catalog Number:Observed MW:AG2842422-25 kDa

**Applications** 

Tested Applications:

WB, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity:

human
Cited Species:
human

Positive Controls:

WB: HL-60 cells, human placenta tissue, THP-1 cells, U-

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

CloneNo.:

2G12E6

937 cells

**Background Information** 

PYCARD, also known as ASC or TMS1, is a 195 amino acid protein containing an N-terminal Pyrin-like domain (PYD) and an 87 residue C-terminal Caspase-associated recruitment domain (CARD). It promotes caspase-mediated apoptosis which is mediated predominantly through the activation of caspase-9.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Jianyang Liu	34733914	Ann Transl Med	WB
Yiping Leng	32912629	Biochem Biophys Res Commun	WB
Yingchun Zhao	34663844	Sci Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**





Various lysates were subjected to SDS PAGE followed by western blot with 67494-1-1g (ASC/TMS1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.

human placenta tissue were subjected to SDS PAGE followed by western blot with 67494-1-lg (ASC/TMS1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.