

For Research Use Only

MRE11A Monoclonal antibody

Catalog Number: 67570-1-Ig



Basic Information

Catalog Number: 67570-1-Ig	GenBank Accession Number: BC063458	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4361	CloneNo.: 1B12B2
Source: Mouse	UNIPROT ID: P49959	Recommended Dilutions: WB 1:3000-1:50000
Isotype: IgG1	Full Name: MRE11 meiotic recombination 11 homolog A (S. cerevisiae)	
Immunogen Catalog Number: AG28847	Observed MW: 81 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, 4T1 cells
Species Specificity: Human, Mouse, Rat	

Background Information

Mre11 (meiotic recombination 11), also named as Mre11A, is a component of the MRN(MRE11/RAD50/NBS1) complex which is a versatile complex, playing a key role in the sensing, processing, and repair of DSBs. MRE11 possesses both endonuclease and 3'-5' exonuclease activity. It contains two DNA-binding domains (DBDs), enabling it to bind both single-stranded DNA (ssDNA) and double-stranded DNA (dsDNA). After DSBs, MRE11 is loaded onto DSBs sites and cleaves DNA by cooperating with CtIP (CtBP [C terminal binding protein] interacting protein) to initiate end resection. It's reported that MRE11 is over-expressed in breast cancers. (PMID:38128537; 22914783)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67570-1-Ig (MRE11A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.