For Research Use Only

CCDC6 Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

1D6A8

Catalog Number: 67637-1-PBS

Basic Information

Catalog Number:

67637-1-PBS

100ug, Concentration: 1 mg/ml by Nanodrop:

Mouse Isotype:

lgG1

Immunogen Catalog Number: AG6952

UNIPROT ID: Q16204 Full Name:

GeneID (NCBI):

BC036757

coiled-coil domain containing 6

GenBank Accession Number:

Calculated MW: 474 aa, 53 kDa Observed MW:

65 kDa

Applications

Tested Applications: WB, IF, Indirect ELISA

Species Specificity:

Human, mouse, rat

Background Information


CCDC6 (Coiled-coil domain-containing protein 6) is also named as Protein H4, D10S170 and TST1. CCDC6, was initially isolated as part of a tumorigenic DNA originated by the fusion of CCDC6 with the tyrosine kinase of RET receptor. CCDC6 has been considered as an accidental partner of the RET protooncogene, providing the promoter and the first 101 aa necessary for the constitutive activation of the oncogenic Tyrosine Kinase (TK) RET in thyroid cells. The 65 kDa product of CCDC6 has a nuclear transfer sequence with no transmembrane domains and is predicted to locate in both the nucleus and the cytoplasm (PMID: 29044514). The CCDC6 is a phosphoprotein, predicted target of several S/T kinases which can modulate the protein stability and the intracellular shuttling into the nucleus upon different cellular signals mediated by ERK1/2, ATM and CDK1/2. CCDC6 is involved in cellular response to DNA damage mediated by ATM, with the final result of promoting cellular apoptosis. CCDC6 depleted cells are considered defective of DNA repair checkpoint and proceed faster than the control cells in the cell cycle upon induced DNA damage (PMID: 22655027).

Storage


Store at -80°C. Storage Buffer: PBS Only

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67637-1-lg (CCDC6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67637-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 67637-1-lg (CCDC6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67637-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using CCDC6 antibody (67637-1-Ig, Clone: 1D6A8) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 67637-1-PBS in a different storage buffer formulation.