For Research Use Only

VAMP2 Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

1G7E8

Catalog Number: 67822-1-PBS

Basic Information

Catalog Number: GenBank Accession Number:

67822-1-PBS BC002737

GeneID (NCBI): 100ug, Concentration: 1 mg/ml by

Nanodrop: **UNIPROT ID:** P63027 Mouse Full Name:

Isotype: vesicle-associated membrane protein

IgG3 2 (synaptobrevin 2) Immunogen Catalog Number: Calculated MW: AG17908 13 kDa

> Observed MW: 19 kDa

Applications

Tested Applications: WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, Mouse, Rat, Pig, Rabbit

Background Information

VAMP2 (vesicle-associated membrane protein 2), also named as synaptobrevin 2, is a member of the SNARE (soluble NSF-attachment protein receptor) family proteins. Characterized by a common sequence called the SNARE motif, SNARE proteins are involved in membrane fusion and vesicular transport (PMID: 11252968). VAMP2, with a molecular mass of 15-19 kDa, consists of a short N-terminal sequence, a SNARE motif, and a C-terminal transmembrane region. It is required for fast calcium-triggered synaptic vesicle fusion. VAMP2 forms a stable complex with STX1 (syntaxin 1) and SNAP25 (synaptosomal-associated protein 25) during synaptic vesicle fusion $(PMID: 16793874). \ It also forms a distinct complex with synaptophysin. \ VAMP2 is expressed in nervous system and a synaptophysin of the complex with synaptophysin of the complex with synaptophysin. \ VAMP2 is expressed in nervous system and the complex with synaptophysin of the complex with synaptophysin o$ some non-neuronal tissues, such as skeletal muscle (PMID: 18570252).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67822-1-lg (VAMP2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67822-1-Ig (VAMP2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67822-1-Ig (VAMP2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed U-87 MG cells using VAMP2 antibody (67822-1-Ig, Clone: 1G7E8) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.