

For Research Use Only

# Phospho-MEK1 (Thr386) Monoclonal antibody, PBS Only (Detector)

Catalog Number: 68015-1-PBS



## Basic Information

<b>Catalog Number:</b> 68015-1-PBS	<b>GenBank Accession Number:</b> BC139729	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ug, Concentration: 1mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5604	<b>CloneNo.:</b> 1G6A2
<b>Source:</b> Mouse	<b>ENSEMBL Gene ID:</b> ENSG00000169032	
<b>Isotype:</b> IgG1	<b>UNIPROT ID:</b> Q02750	
	<b>Full Name:</b> mitogen-activated protein kinase kinase 1	
	<b>Calculated MW:</b> 43 kDa	
	<b>Observed MW:</b> 40-50 kDa	

## Applications

**Tested Applications:**  
WB, IF/ICC, Cytometric bead array, Indirect ELISA

**Species Specificity:**  
human, mouse

## Product Information

68015-1-PBS targets Phospho-MEK1 (Thr386) as part of a matched antibody pair.

MP50180-1: 67872-1-PBS capture and 68015-1-PBS detection (validated in Cytometric bead array)

Unconjugated mouse monoclonal antibody pair in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.

## Background Information

MAP2K1 encodes MAPK1, also known as MEK1. MEK1 variants can enhance MEK1 expression and ERK1 phosphorylation that together lead to continuous activation of MEK/ERK signaling pathway. MEK1 bind directly to ERK2 through a region in the N terminus of MEK. In addition, a proline-rich (PR) regulatory sequence in MEK is also involved in MEK-ERK association and signal propagation. The coupling between MEK1 and ERK2 is enhanced through phosphorylation on S298 in the MEK1 PR region, whereas phosphorylation on MEK1 T292 releases the complex. MEK1 T292 is a substrate of ERK2, but the site is also phosphorylated at a basal level when ERK2 is inhibited, suggesting several regulators of this site. Although the S298 site in MEK2 has been conserved, it lacks the T292 phosphorylation site, and it is not a substrate of PAK1. (PMID: 31972311, PMID: 17928366, PMID: 22177953)

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Non-treated NIH/3T3 cells and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 68015-1-Ig (Phospho-MEK1 (Thr386) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control. This data was developed using the same antibody clone with 68015-1-PBS in a different storage buffer formulation.




Non-treated HeLa cells and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 68015-1-Ig (Phospho-MEK1 (Thr386) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control. This data was developed using the same antibody clone with 68015-1-PBS in a different storage buffer formulation.



Non-treated HeLa cells, phosphatase inhibitor treated and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 68015-1-Ig (Phospho-MEK1 (Thr386) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68015-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Calyculin A treated HeLa cells using Phospho-MEK1 (Thr386) antibody (68015-1-Ig, Clone: 1G6A2 ) at dilution of 1:400 and Multi-rAb CoralLite® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002). This data was developed using the same antibody clone with 68015-1-PBS in a different storage buffer formulation.



Cytometric bead array in cell lysate using MP50180-1, Phospho-MEK1 (Thr386) Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 67872-1-PBS. Detection antibody: 68015-1-PBS. Cell lysate: Non-treated HeLa and Calyculin A treated HeLa (30µg/well). Non-related target OAT Monoclonal Matched Antibody Pair (MP50109-1P) was served as control.