For Research Use Only

## TMPRSS2 Monoclonal antibody, PBS Only



Catalog Number: 68264-1-PBS

**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

68264-1-PBS

GeneID (NCBI):

Protein G purification

CloneNo.:

100ug, Concentration: 1 mg/ml by Nanodrop:

**UNIPROT ID:** 

BC051839

1H7E9

015393

Mouse Isotype:

Full Name: transmembrane protease, serine 2

lgG1

Calculated MW:

Immunogen Catalog Number:

54 kDa

AG5824

Observed MW:

50 kDa

**Applications** 

**Tested Applications:** 

WB, Indirect ELISA

Species Specificity:

Human, Pig

## **Background Information**

TMPRSS2, also named as PRSS10, is a type II transmembrane serine protease which is highly expressed by the epithelium of the human prostate gland. TMPRSS2 may contribute to prostate tumour metastasis via the activation of PAR-2. TMPRSS2 is a Serine protease that proteolytically cleaves and activates the viral spike glycoproteins which facilitate virus-cell membrane fusions. TMPRSS2 is a host cell factor that is critical for the spread of several clinically relevant viruses, including influenza A viruses and coronaviruses (PMID: 23468491, 30626688). SARS-CoV-2 uses the SARS-CoV receptor ACE2 for entry and the serine protease TMPRSS2 for S protein priming. The initial spike protein priming by TMPRSS2 is essential for the entry and viral spread of SARS-CoV-2 through interaction with the ACE2 receptor (PMID: 32142651, 30626688). Camostat mesylate, an inhibitor of TMPRSS2, can block SARS-CoV-2 infection of lung cells (PMID: 32142651). The MW of TMPRSS2 is about 65-70 kDa. It can be cleaved into some chains with MW 54 kDa, 31 kDa and 26 kDa (PMID: 25734995, 20382709, 26018085).


Storage


Storage:

Store at -80°C. Storage Buffer:

PBS Only


## **Selected Validation Data**







pig kidney tissue were subjected to SDS PAGE followed by western blot with 68264-1-lg (TMPRSS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68264-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68264-1-lg (TMPRSS2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68264-1-PBS in a different storage buffer formulation.