For Research Use Only

GATC Monoclonal antibody

Catalog Number: 68405-1-lg



Basic Information

Catalog Number: GenBank Accession Number:

68405-1-lg BC107145 Protein A purification

GeneID (NCBI): CloneNo.: Size: 2F9C8 150ul , Concentration: 1000 $\mu g/ml$ by 283459

Nanodrop; **UNIPROT ID:** Recommended Dilutions: WB 1:2000-1:10000 Source: 043716 IF/ICC 1:200-1:800 Mouse Full Name:

Isotype: glutamyl-tRNA(Gln)

amidotransferase, subunit C homolog lgG2b

(bacterial) Immunogen Catalog Number: AG19857 Calculated MW:

136 aa, 15 kDa Observed MW:

15 kDa

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

human

Positive Controls:

WB: LNCaP cells, A549 cells, HEK-293 cells, K-562 cells

Purification Method:

IF/ICC: HEK-293 cells,

Background Information

GATC encodes a subunit of the glutamyl-tRNA(Gln) amidotransferase protein complex (GatCAB), which plays a role in aminoacylation of mitochondrial glutaminyl mt-tRNA. The complex plays an role in mitochondrial translation and protein synthesis (PubMed: 30283131).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1%BSA

Selected Validation Data








1x10^6 HEK-293 cells were intracellularly stained with 0.25 ug GATC Monoclonal antibody (68405-1-1g, Clone:2F9C8) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (5A00013-1)(red), or 0.25 ug Mouse IgG2b isotype control Mouse McAb (66360-3-Ig, Clone: 11B8C4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

A549 cells were subjected to SDS PAGE followed by western blot with 68405-1-lg (GATC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

Various lysates were subjected to SDS PAGE followed by western blot with 68405-1-Ig (GATC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using GATC antibody (68405-1-Ig, Clone: 2F9C8) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1).