### For Research Use Only

# Hexokinase 1 Monoclonal antibody

Catalog Number: 68419-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

68419-1-lg BC008730 GeneID (NCBI):

150ul, Concentration: 1000 µg/ml by 3098 2B4F9

Full Name: Source: hexokinase 1

Mouse Calculated MW:

Isotype: 102 kDa IgG2a Observed MW:

Immunogen Catalog Number: 100 kDa

**Tested Applications:** 

IHC, IP, WB, ELISA Species Specificity:

**Purification Method:** Protein A purification

CloneNo.:

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for IP and 1:1000-1:4000

for WB

IHC 1:500-1:2000

**Applications** 

Human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: HeLa cells, HT-1080 cells, U-87 MG cells, SK-BR-3 cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, pig brain tissue

IP: HeLa cells,

IHC: mouse brain tissue,

## **Background Information**

The hexokinases (HKs) catalyze the first obligatory step in glucose metabolism to generate glucose-6-phosphate. This not only furthers glucose entry by maintaining the concentration gradient for facilitated glucose influx but also provides the first intermediate for essentially all major pathways using glucose. There are four conventional HK isoforms, HK1/2/3/4, encoded by four different genes. Most adult tissues express only HK1. Muscle and adipose tissue use HK2 for glycolysis, liver, and pancreatic beta cells express HK4 (also called glucokinase) and do not express HK1 or HK2. (PMID: 31434645, PMID: 31848318)

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68419-1-1g (HK1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-Hexokinase 1(IP:68419-1-Ig, 4ug; Detection:68419-1-Ig 1:2000) with HeLa cells lysate 1560 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 68419-1-Ig (Hexokinase 1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).