For Research Use Only

STEAP4 Monoclonal antibody, PBS Only

Catalog Number:68465-1-PBS


Antibodies | ELISA kits | Proteins WWW.ptglab.com


Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 68465-1-PBS BC020600 Protein A purification GenelD (NCBI): CloneNo.: Size: 100ug , Concentration: 1mg/ml by 79689 2A12G6 Nanodrop: UNIPROT ID: Source: Q687X5 Mouse Full Name: Isotype: STEAP family member 4 lgG1 Calculated MW: Immunogen Catalog Number: 52 kDa AG30137 **Observed MW:** 52 kDa **Applications Tested Applications:** WB, ELISA **Species Specificity:** human, mouse, rat **Background Information** STEAP4 (also called STAMP2), is a member of the STEAP family. Structurally, It is a 459 amino acid protein characterized by a molecular topology of six transmembrane domains. The cytoplasmic N-terminal domain of STEAP4 shows structural similarity with bacterial and archaeal FNO oxidoreductase and STEAP4 exhibits a strong iron reductase activity. Studies in mice and human suggest that STEAP4 maybe involved in adipocyte development and metabolism, and may contribute to the normal biology of the prostate cell, as well as prostate cancer progression. Storage Storage: Store at -80°C. Storage Buffer: PBS Only

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 68465-1-Ig (STEAP4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 68465-1-PBS in a different storage buffer formulation. Various lysates were subjected to SDS PAGE followed by western blot with 68465-1-1g (STEAP4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68465-1-PBS in a different storage buffer formulation.